

Stabilizers of functional Menger systems

Wiesław A. Dudek and Valentin S. Trokhimenko

Abstract

A functional Menger system is a set of n -place functions containing n projections and closed under the so-called Menger's composition of n -place functions. We give the abstract characterization for subsets of these functional systems which contain functions having one common fixed point.

1 Introduction

Investigation of multiplace functions by algebraic methods plays a very important role in modern mathematics where we consider various operations on sets of functions, which are naturally defined. The basic operation for functions is superposition (composition), but there are some other naturally defined operations, which are also worth of consideration. For example, the operation of set-theoretic intersection and the operation of projections (see for example [1, 2, 3, 6, 7]). The central role in these study play sets of functions with fixed points. The study of such sets for functions of one variable was initiated by B. M. Schein in [4] and [5]. Next, for sets of functions of n variables, was continued by V.S. Trokhimenko (see [8, 9]).

In this paper, we consider the sets of n -place functions containing n -projections and closed under the so-called Menger's composition of n -place functions. For such functional systems we give the abstract characterization for subsets of functions having one common fixed point.

2 Preliminaries

Let A^n be the n -th Cartesian product of a set A . Any partial map from A^n into A is called an n -place function on A . The set of all such maps is denoted by $\mathcal{F}(A^n, A)$. On $\mathcal{F}(A^n, A)$ we define one $(n+1)$ -ary superposition $\mathcal{O}: (f, g_1, \dots, g_n) \mapsto f[g_1 \dots g_n]$, called the *Menger's composition*, and n unary operations $\mathcal{R}_i: f \mapsto \mathcal{R}_i f$, $i \in \overline{1}, n = \{1, \dots, n\}$ putting

$$f[g_1 \dots g_n](a_1, \dots, a_n) = f(g_1(a_1, \dots, a_n), \dots, g_n(a_1, \dots, a_n)), \quad (1)$$

$$\mathcal{R}_i f(a_1, \dots, a_n) = a_i, \text{ where } (a_1, \dots, a_n) \in \text{pr}_1 f \quad (2)$$

for $f, g_1, \dots, g_n \in \mathcal{F}(A^n, A)$, $(a_1, \dots, a_n) \in A^n$, where $\text{pr}_1 f$ denotes the domain of a function f . It is assumed the left and right hand side of equality (1) are defined, or not defined, simultaneously. Algebras of the form $(\Phi, \mathcal{O}, \mathcal{R}_1, \dots, \mathcal{R}_n)$,

where $\Phi \subset \mathcal{F}(A^n, A)$, are called *functional Menger systems of n -place functions*. Algebras of the form $(\Phi, \mathcal{O}, \cap, \mathcal{R}_1, \dots, \mathcal{R}_n)$, where $\cap$ is a set-theoretic intersection, are called *functional Menger $\cap$ -algebras of n -place functions*. In the literature such algebras are also called *functional Menger $\mathcal{P}$ -algebras* (see [1] and [9]).

Let a be some fixed element (point) of A . The *stabilizer of a* is the set H_Φ^a of such functions from Φ for which a is a fixed point, i.e., the set

$$H_\Phi^a = \{f \in \Phi \mid f(a, \dots, a) = a\}.$$

Let (G, o) be a nonempty set with one $(n+1)$ -ary operation

$$o: (x_0, x_1, \dots, x_n) \mapsto x_0[x_1 \dots x_n].$$

An abstract algebra $\mathcal{G} = (G, o, R_1, \dots, R_n)$ of type $(n+1, 1, \dots, 1)$ for all $i, k \in \overline{1, n}$ satisfying the following axioms:

$$\begin{aligned} \mathbf{A}_1: & \quad x[y_1 \dots y_n][z_1 \dots z_n] = x[y_1[z_1 \dots z_n] \dots y_n[z_1 \dots z_n]], \\ \mathbf{A}_2: & \quad x[R_1 x \dots R_n x] = x, \\ \mathbf{A}_3: & \quad x[\bar{u} \mid_i z][R_1 y \dots R_n y] = x[\bar{u} \mid_i z][R_1 y \dots R_n y], \\ \mathbf{A}_4: & \quad R_i x[R_1 y \dots R_n y] = (R_i x)[R_1 y \dots R_n y], \\ \mathbf{A}_5: & \quad x[R_1 y \dots R_n y][R_1 z \dots R_n z] = x[R_1 z \dots R_n z][R_1 y \dots R_n y], \\ \mathbf{A}_6: & \quad R_i x[y_1 \dots y_n] = R_i(R_k x)[y_1 \dots y_n], \\ \mathbf{A}_7: & \quad (R_i x)[y_1 \dots y_n] = y_i[R_1(x[y_1 \dots y_n]) \dots R_n(x[y_1 \dots y_n])], \end{aligned}$$

where $x[\bar{u} \mid_i z]$ means $x[u_1 \dots u_{i-1} z u_{i+1} \dots u_n]$, is called a *functional Menger system of rank n* .

An algebra $\mathcal{G}_\wedge = (G, o, \wedge, R_1, \dots, R_n)$ of type $(n+1, 2, 1, \dots, 1)$, where $(G, o, R_1, \dots, R_n)$ is a functional Menger system of rank n and $(G, \wedge)$ is a semi-lattice, is called a *functional Menger $\wedge$ -algebra of rank n* if it satisfies the identities:

$$\begin{aligned} \mathbf{A}_8: & \quad x \wedge y[R_1 z \dots R_n z] = (x \wedge y)[R_1 z \dots R_n z], \\ \mathbf{A}_9: & \quad x \wedge y = x[R_1(x \wedge y) \dots R_n(x \wedge y)], \\ \mathbf{A}_{10}: & \quad (x \wedge y)[z_1 \dots z_n] = x[z_1 \dots z_n] \wedge y[z_1 \dots z_n]. \end{aligned}$$

Any Menger algebra of rank n , i.e., an abstract groupoid (G, o) with an $(n+1)$ -ary operation satisfying $\mathbf{A}_1$ is isomorphic to some set of n -place functions closed under Menger's composition [6]. Functional Menger $\wedge$ -algebras and Menger systems of rank n are isomorphic, respectively, to some functional Menger $\cap$ -algebras and Menger systems of n -place functions (see [1] and [7]). Each homomorphism of such abstract algebras into corresponding algebras of n -place functions is called a *representation by n -place functions*. Representations which are isomorphisms are called *faithful*.

Let $(P_i)_{i \in I}$ be the family of representations of a Menger algebra (G, o) of rank n by n -place functions defined on sets $(A_i)_{i \in I}$, respectively. By the *union*

of this family we mean the map $P: g \mapsto P(g)$, where $g \in G$, and $P(g)$ is an n -place function on $A = \bigcup_{i \in I} A_i$ defined by

$$P(g) = \bigcup_{i \in I} P_i(g).$$

If $A_i \cap A_j = \emptyset$ for all $i, j \in I$, $i \neq j$, then P is called the *sum* of $(P_i)_{i \in I}$ and is denoted by $P = \sum_{i \in I} P_i$. It is not difficult to see that the sum of representations is a representation, but the union of representations may not be a representation (see for example [1 – 7]).

For every representation $P: G \rightarrow \mathcal{F}(A^n, A)$ of an algebra (G, o) and every element $a \in G$ by H_P^a we denote the set of elements of G corresponding to these n -place functions for which a is a fixed point, i.e.,

$$H_P^a = \{g \in G \mid P(g)(a, \dots, a) = a\}.$$

Let $\mathcal{G}$ be a functional Menger system of rank n , x – an individual variable. By $T_n(G)$ we denote the set of transformations $t: x \mapsto t(x)$ on G such that:

- (a) $x \in T_n(G)$,
- (b) if $t(x) \in T_n(G)$, then $a[\bar{b} \mid_i t(x)] \in T_n(G)$ and $R_i t(x) \in T_n(G)$ for all $a \in G$, $\bar{b} \in G^n$ and $i \in \overline{1, n}$,
- (c) $T_n(G)$ contains only elements determined in (a) and (b).

Let us remind that a nonempty subset H of G is called

- *quasi-stable*, if for all $x \in G$

$$x \in H \longrightarrow x[x \dots x] \in H,$$

- $\wedge$ -*quasi-stable*, if for all $x \in G$

$$x \in H \longrightarrow x[x \dots x] \wedge x \in H,$$

- *stable*, if for all $x, y_1, \dots, y_n \in G$

$$x, y_1, \dots, y_n \in H \longrightarrow x[y_1 \dots y_n] \in H,$$

- $\wedge$ -*stable*, if for all $x, y \in G$

$$x, y \in H \longrightarrow x \wedge y \in H,$$

- *l -unitary*, if for every $x, y \in G$

$$x[y \dots y] \in H \wedge y \in H \longrightarrow x \in H,$$

- *v-unitary*, if for all $x, y_1, \dots, y_n \in G$

$$x[y_1 \dots y_n] \in H \wedge y_1, \dots, y_n \in H \longrightarrow x \in H,$$

- a *normal v-complex*, if for all $x, y \in G, t \in T_n(G)$

$$x, y \in H \wedge t(x) \in H \longrightarrow t(y) \in H,$$

- an *l-ideal*, if for all $x, y_1, \dots, y_n \in G$

$$(y_1, \dots, y_n) \in G^n \setminus (G \setminus H)^n \longrightarrow x[y_1 \dots y_n] \in H.$$

A binary relation $\rho \subset G \times G$ is called

- *stable*, if

$$(x, y), (x_1, y_1), \dots, (x_n, y_n) \in \rho \longrightarrow (x[x_1 \dots x_n], y[y_1 \dots y_n]) \in \rho$$

for all $x, y, x_i, y_i \in G, i \in \overline{1, n}$,

- *l-regular*, if

$$(x, y) \in \rho \longrightarrow (x[z_1 \dots z_n], y[z_1 \dots z_n]) \in \rho$$

for all $x, y, z_i \in G, i \in \overline{1, n}$,

- *v-regular*, if

$$(x_1, y_1), \dots, (x_n, y_n) \in \rho \longrightarrow (z[x_1 \dots x_n], z[y_1 \dots y_n]) \in \rho$$

for all $x_i, y_i, z \in G, i \in \overline{1, n}$,

- *i-regular*, where $i \in \overline{1, n}$, if

$$(x, y) \in \rho \longrightarrow (u[\bar{w} \mid_i x], u[\bar{w} \mid_i y]) \in \rho$$

for all $x, y, u \in G, \bar{w} \in G^n$,

- *v-negative*, if

$$(x[y_1 \dots y_n], y_i) \in \rho$$

for all $x, y_1, \dots, y_n \in G$ and $i \in \overline{1, n}$.

On $\mathcal{G}$ we define two binary relations ζ and χ putting

$$(x, y) \in \zeta \longleftrightarrow x = y[R_1 x \dots R_n x], \quad (x, y) \in \chi \longleftrightarrow (R_1 x, R_1 y) \in \zeta.$$

The first relation is a stable order, the second is an *l-regular* and *v-negative* quasi-order containing ζ (see [7]). For these two relations the following conditions are valid:

$$\begin{aligned} x \leq y &\longrightarrow R_i x \leq R_i y, \quad i \in \overline{1, n}, & x \sqsubset y &\longleftrightarrow R_i x \leq R_i y, \quad i \in \overline{1, n}, \\ x \sqsubset y &\longleftrightarrow x[R_1 y \dots R_n y] = x, & (R_i x)[y_1 \dots y_n] &\leq y_i, \quad i \in \overline{1, n}, \\ x[R_1 y_1 \dots R_n y_n] &\leq x, & R_i x &= R_i R_k x, \quad i, k \in \overline{1, n}, \end{aligned}$$

where $x \leq y \iff (x, y) \in \zeta$, and $x \sqsubset y \iff (x, y) \in \chi$.

Let W be the empty set or an l -ideal which is an $\mathcal{E}$ -class of a v -regular equivalence relation $\mathcal{E}$ defined on a Menger algebra (G, o) of rank n . Denote by $(H_a)_{a \in A_{\mathcal{E}}}$ the family of all $\mathcal{E}$ -classes (uniquely indexed by elements of some set $A_{\mathcal{E}}$) such that $H_a \neq W$. Next, for every $g \in G$ we define on $A_{\mathcal{E}}$ an n -place function $P_{(\mathcal{E}, W)}(g)$ putting

$$P_{(\mathcal{E}, W)}(g)(a_1, \dots, a_n) = b \iff g[H_{a_1} \dots H_{a_n}] \subset H_b, \quad (3)$$

where $(a_1, \dots, a_n) \in \text{pr}_1 P_{(\mathcal{E}, W)}(g) \iff g[H_{a_1} \dots H_{a_n}] \cap W = \emptyset$, and H_b is an $\mathcal{E}$ -class containing all elements of the form $g[h_1 \dots h_n]$, $h_i \in H_{a_i}$, $i \in \overline{1, n}$. It is not difficult to see that the map $P_{(\mathcal{E}, W)}: g \mapsto P_{(\mathcal{E}, W)}(g)$ satisfies the identity

$$P_{(\mathcal{E}, W)}(g[g_1 \dots g_n]) = P_{(\mathcal{E}, W)}(g)[P_{(\mathcal{E}, W)}(g_1) \dots P_{(\mathcal{E}, W)}(g_n)], \quad (4)$$

i.e., $P_{(\mathcal{E}, W)}$ is a representation of (G, o) by n -place functions. This representation will be called *simplest*.

3 Stabilizers

Definition 1. A nonempty subset H of G is called a *stabilizer* of a functional Menger system $\mathcal{G}$ (or a functional Menger λ -algebra $\mathcal{G}_{\lambda} = (G, \lambda, o, R_1, \dots, R_n)$) of rank n if there exists a representation P of $\mathcal{G}$ (respectively, $\mathcal{G}_{\lambda}$) by n -place functions on some set A , such that $H = H_P^a$ for some point $a \in A$ common for all elements from H .

Theorem 1. For a nonempty subset H of G to be a stabilizer of a functional Menger system $\mathcal{G}$ of rank n , it is necessary and sufficient to be a quasi-stable l -unitary normal v -complex contained in some subset U of G such that $R_i U \subset H$, $R_i(G \setminus U) \subset G \setminus U$ and

$$x \in H \wedge y \in H \wedge t(x) \in U \implies t(y) \in U, \quad (5)$$

$$x = y[R_1 x \dots R_n x] \in U \wedge u[\bar{w} |_i y] \in H \implies u[\bar{w} |_i x] \in H, \quad (6)$$

$$x = y[R_1 x \dots R_n x] \in U \wedge u[\bar{w} |_i y] \in U \implies u[\bar{w} |_i x] \in U \quad (7)$$

for all $x, y \in G$, $\bar{w} \in G^n$, $t \in T_n(G)$ and $i \in \overline{1, n}$, where the symbol $u[\bar{w} |_i]$ may be empty.¹

Proof. Necessity. Let H_{Φ}^a be a stabilizer of a for a functional Menger system $(\Phi, \mathcal{O}, \mathcal{R}_1, \dots, \mathcal{R}_n)$ of n -place functions. If $f \in H_{\Phi}^a$, i.e., $f(a, \dots, a) = a$, then

$$f[f \dots f](a, \dots, a) = f(f(a, \dots, a), \dots, f(a, \dots, a)) = f(a, \dots, a) = a.$$

Thus $f[f \dots f] \in H_{\Phi}^a$. This means that H_{Φ}^a is quasi-stable.

¹If $u[\bar{w} |_i]$ is the empty symbol, then $u[\bar{w} |_i x]$ is equal to x .

Since, for $f[g \dots g] \in H_\Phi^a$ and $g \in H_\Phi^a$ we have

$$a = f[g \dots g](a, \dots, a) = f(g(a, \dots, a), \dots, g(a, \dots, a)) = f(a, \dots, a),$$

then $f \in H_\Phi^a$, therefore H_Φ^a is l -unitary.

Moreover, if $f(\bar{a}) = g(\bar{a})$ for some $f, g \in \Phi$, where $\bar{a} = (a_1, \dots, a_n)$, then, as it is not difficult to see, $t(f)(\bar{a}) = t(g)(\bar{a})$ for every $t \in T_n(\Phi)$. So, $f, g, t(f) \in H_\Phi^a$ implies $t(g) \in H_\Phi^a$. Therefore H_Φ^a is a normal v -complex.

It is clear that $H_\Phi^a \subset U_\Phi^a = \{f \in \Phi \mid (a, \dots, a) \in \text{pr}_1 f\}$, $\mathcal{R}_i U_\Phi^a \subset H_\Phi^a$ and $\mathcal{R}_i(\Phi \setminus U_\Phi^a) \subset \Phi \setminus U_\Phi^a$. If $t(f) \in U_\Phi^a$ for some $f, g \in H_\Phi^a$ and $t \in T_n(\Phi)$, then $(a, \dots, a) \in \text{pr}_1 t(f)$, which, together with $f(a, \dots, a) = a = g(a, \dots, a)$, gives $t(f)(a, \dots, a) = t(g)(a, \dots, a)$. Therefore, $t(g) \in U_\Phi^a$. So, the condition (5) is satisfied.

To prove (6) assume $f = g[\mathcal{R}_1 f \dots \mathcal{R}_n f] \in U_\Phi^a$, i.e., $f \subset g$ and $f \in U_\Phi^a$. This implies $(a, \dots, a) \in \text{pr}_1 f$ and $f(a, \dots, a) = g(a, \dots, a)$. So, for $\alpha[\bar{\chi} |_i g] \in H_\Phi^a$, where $\alpha \in \Phi$ and $\bar{\chi} = (\chi_1, \dots, \chi_n) \in \Phi^n$, we have

$$\begin{aligned} a &= \alpha[\bar{\chi} |_i g](a, \dots, a) = \alpha(\bar{\chi}(a, \dots, a) |_i g(a, \dots, a)) \\ &= \alpha(\bar{\chi}(a, \dots, a) |_i f(a, \dots, a)) = \alpha[\bar{\chi} |_i f](a, \dots, a), \end{aligned}$$

where $\bar{\chi}(a, \dots, a)$ is $\chi_1(a, \dots, a), \dots, \chi_n(a, \dots, a)$. Thus $\alpha[\bar{\chi} |_i f] \in H_\Phi^a$, which completes the proof of (6). The proof of (7) is analogous.

Sufficiency. Let H and U be two subsets of G satisfying all the conditions of the theorem. First we shall prove the following implications:

$$x \leq y \wedge x \in H \longrightarrow y \in H, \quad (8)$$

$$x \sqsubset y \wedge x \in U \longrightarrow y \in U. \quad (9)$$

Indeed, $x \leq y$ means $x = y[R_1 x \dots R_n x]$. Since $x \in H$, $H \subset U$ and $R_i U \subset H$, we have $R_i x \in H$ for every $i \in \overline{1, n}$. This, together with the fact that H is an l -unitary normal v -complex, implies that H is v -unitary. So, $y[R_1 x \dots R_n x] \in H$ and $R_i x \in H$ for every $i \in \overline{1, n}$, which by the v -unitarity of H gives $y \in H$. This proves (8).

Now, if $x \sqsubset y$ and $x \in U$, then $R_1 x \leq R_1 y$, i.e., $R_1 x = (R_1 y)[R_1 x \dots R_n x]$. From $R_i U \subset H$ it follows $R_i x \in H$, so, applying the v -unitarity of H to $(R_1 y)[R_1 x \dots R_n x] \in H$ we obtain $R_1 y \in H$, whence we get $R_1 y \in U$. Since $R_i(G \setminus U) \subset G \setminus U$ means that

$$R_i x \in U \longrightarrow x \in U \quad (10)$$

for every $x \in G$ and $i \in \overline{1, n}$, from $R_1 y \in U$ it follows $y \in U$. This completes the proof of (9).

The set $G \setminus U$ is an l -ideal of $\mathcal{G}$. In fact, the v -negativity of χ and $x \in G \setminus U$ imply $u[\bar{w} |_i x] \sqsubset x$, whence, according to (9), we conclude $u[\bar{w} |_i x] \in G \setminus U$ for all $i \in \overline{1, n}$, $u \in G$ and $\bar{w} \in G^n$. So, $G \setminus U$ is an l -ideal. Using this fact it is easy

to show that

$$x \leq y \wedge x \in U \wedge t(y) \in H \longrightarrow t(x) \in H, \quad (11)$$

$$x \leq y \wedge x \in U \wedge t(y) \in U \longrightarrow t(x) \in U \quad (12)$$

for all $x, y \in G$ and $t \in T_n(G)$.

On G we define two binary relations $\mathcal{E}_H$ and $\mathcal{E}_U$ putting

$$\begin{aligned} \mathcal{E}_H &= \left\{ (x, y) \mid (\forall t \in T_n(G)) (t(x) \in H \longleftrightarrow t(y) \in H) \right\}, \\ \mathcal{E}_U &= \left\{ (x, y) \mid (\forall t \in T_n(G)) (t(x) \in U \longleftrightarrow t(y) \in U) \right\}. \end{aligned}$$

These relations are v -regular equivalences. $\mathcal{E} = \mathcal{E}_H \cap \mathcal{E}_U$ also is a v -regular equivalence. For this relation we have

$$x[\mathcal{E}\langle y_1 \rangle \dots \mathcal{E}\langle y_n \rangle] \subset \mathcal{E}\langle x[y_1 \dots y_n] \rangle \quad (13)$$

for all $x, y_1, \dots, y_n \in G$, where $\mathcal{E}\langle y_i \rangle$ denotes an equivalence class of $\mathcal{E}$ containing y_i . Moreover, $G \setminus U$ is an equivalence class of $\mathcal{E}$.

Also H is an $\mathcal{E}$ -class. To prove this fact it is sufficient to verify the following two conditions:

$$g_1 \in H \wedge g_2 \in H \longrightarrow g_1 \equiv g_2(\mathcal{E}), \quad (14)$$

$$g_1 \equiv g_2(\mathcal{E}) \wedge g_1 \in H \longrightarrow g_2 \in H. \quad (15)$$

Let $g_1, g_2 \in H$ and $t(g_1) \in U$ for some $t \in T_n(G)$. Then, from (5), it follows $t(g_2) \in U$. Similarly, from $g_1, g_2 \in H$ and $t(g_2) \in U$, we conclude $t(g_1) \in U$. Hence, $g_1 \equiv g_2(\mathcal{E}_U)$. If $g_1, g_2, t(g_1) \in H$, then, in view of the fact that H is a normal v -complex, we have $t(g_2) \in H$. Similarly, from $g_1, g_2, t(g_2) \in H$ we deduce $t(g_1) \in H$. So, $g_1 \equiv g_2(\mathcal{E}_H)$. Thus $g_1 \equiv g_2(\mathcal{E})$. This proves (14).

Now let $g_1 \in H$ and $g_1 \equiv g_2(\mathcal{E})$. Then $g_1 \equiv g_2(\mathcal{E}_H)$, which means that for all $t \in T_n(G)$ we have $t(g_1) \in H \longleftrightarrow t(g_2) \in H$, whence $g_1 \in H \longleftrightarrow g_2 \in H$. So, $g_2 \in H$, i.e., (15) is proved. Consequently, H is an $\mathcal{E}$ -class.

Let $W = G \setminus U$. For every $g \in G$ we consider an n -place function $P_{(\mathcal{E}, W)}(g)$ defined by (3). Since the map $P_{(\mathcal{E}, W)}: g \mapsto P_{(\mathcal{E}, W)}(g)$ satisfies (4), it is a homomorphism with respect to the operation o . It satisfies also the identity

$$P_{(\mathcal{E}, W)}(R_i g) = \mathcal{R}_i P_{(\mathcal{E}, W)}(g). \quad (16)$$

Indeed, for every

$$\bar{a} = (a_1, \dots, a_n) \in \text{pr}_1 P_{(\mathcal{E}, W)}(R_i g),$$

where $H_{a_i} = \mathcal{E}\langle x_i \rangle$, $x_i \in U$, $i \in \overline{1, n}$, we have $(R_i g)[x_1 \dots x_n] \in U$. Whence, according to **A**₇, we obtain $x_i[R_1 g[x_1 \dots x_n] \dots R_n g[x_1 \dots x_n]] \in U$. As $G \setminus U$ is an l -ideal, the last condition implies $R_i g[x_1 \dots x_n] \in U$ for $i \in \overline{1, n}$. Thus $g[x_1 \dots x_n] \in U$, because $g[x_1 \dots x_n] \notin U$ implies $x_i \in G \setminus U$. So, $\bar{a} \in \text{pr}_1 P_{(\mathcal{E}, W)}(g)$. This proves $\text{pr}_1 P_{(\mathcal{E}, W)}(R_i g) \subset \text{pr}_1 P_{(\mathcal{E}, W)}(g)$.

To prove the converse inclusion let $\bar{a} \in \text{pr}_1 P_{(\mathcal{E}, W)}(g)$, where $H_{a_i} = \mathcal{E}\langle x_i \rangle$, $x_i \in U$ for $i \in \overline{1, n}$. Then $g[x_1 \dots x_n] \in U$, which, by $R_k U \subset H \subset U$, gives $R_k g[x_1 \dots x_n] \in U$. Whence, by **A**₆, we get $R_k(R_i)[x_1 \dots x_n] \in U$ for $k, i \in \overline{1, n}$. From this, in view of (10), we deduce $(R_i g)[x_1 \dots x_n] \in U$. Thus $\bar{a} \in \text{pr}_1 P_{(\mathcal{E}, W)}(R_i g)$ for every $i \in \overline{1, n}$.

In this way we have proved

$$\text{pr}_1 P_{(\mathcal{E}, W)}(R_i g) = \text{pr}_1 P_{(\mathcal{E}, W)}(g) = \text{pr}_1 \mathcal{R}_i P_{(\mathcal{E}, W)}(g) \quad (17)$$

for every $g \in G$.

Let $\bar{a} \in \text{pr}_1 P_{(\mathcal{E}, W)}(R_i g)$, i.e., $(R_i g)[x_1 \dots x_n] \in U$, where $H_{a_i} = \mathcal{E}\langle x_i \rangle$, $x_i \in U$, $i \in \overline{1, n}$. Applying the stability of ζ to $(R_i g)[x_1 \dots x_n] \leq x_i$ we obtain $t((R_i g)[x_1 \dots x_n]) \leq t(x_i)$ for every $t \in T_n(G)$ and $i \in \overline{1, n}$. If $t((R_i g)[x_1 \dots x_n]) \in H$, then, according (8), we get $t(x_i) \in H$. For $t(x_i) \in H$, in view of (11), from $(R_i g)[x_1 \dots x_n] \leq x_i$ and $(R_i g)[x_1 \dots x_n] \in U$ we deduce $t((R_i g)[x_1 \dots x_n]) \in H$. So, $(R_i g)[x_1 \dots x_n] \equiv x_i(\mathcal{E}_H)$. Similarly we can prove $(R_i g)[x_1 \dots x_n] \equiv x_i(\mathcal{E}_U)$. Thus $(R_i g)[x_1 \dots x_n] \equiv x_i(\mathcal{E})$ for every $i \in \overline{1, n}$. Therefore $P_{(\mathcal{E}, W)}(R_i g)(a_1, \dots, a_n) = a_i$. Consequently,

$$P_{(\mathcal{E}, W)}(R_i g)(a_1, \dots, a_n) = (\mathcal{R}_i P_{(\mathcal{E}, W)}(g))(a_1, \dots, a_n).$$

This, together with (17), gives (16).

From (4) and (16) it follows that $P_{(\mathcal{E}, W)}$ is a representation of $\mathcal{G}$ by n -place functions.

Observe that

$$g \in H \longleftrightarrow P_{(\mathcal{E}, W)}(g)(a, \dots, a) = a, \quad (18)$$

where a is this element of $A_{\mathcal{E}}$ which is used as index of the $\mathcal{E}$ -class H . In fact, for $g \in H$ the quasi-stability of H implies $g[g \dots g] \in H$. Whence, $g[H \dots H] \subset H$ because H is an $\mathcal{E}$ -class and the relation $\mathcal{E}$ is v -regular. So, $P_{(\mathcal{E}, W)}(g)(a, \dots, a) = a$. Conversely, if $g[H \dots H] \subset H$, then $g[h \dots h] \in H$ for every $h \in H$. From this, by the l -unitarity of H , we get $g \in H$, which completes the proof of (18).

To complete this proof we remind that any algebra $\mathcal{G}$ satisfying the axioms **A**₁ – **A**₇, has a faithful representation by n -place functions [7]. Let P_1 be this representation. Then, as it is not difficult to verify, $P = P_1 + P_{(\mathcal{E}, W)}$ is a faithful representation of $\mathcal{G}$ for which $H = H_P^a$. This completes the proof. $\square$

Let $\mathcal{G}$ be a functional Menger system of rank n and H be some subset of G . We say that a subset X of G is C_H -closed, if for all $a, b, c \in G$, $t \in T_n(G)$ the implication:

$$\left. \begin{array}{l} a = b[R_1 a \dots R_n a] \vee a, b \in H, \\ t(a)[R_1 c \dots R_n c] = t(a), \\ a, t(b) \in X \end{array} \right\} \longrightarrow c \in X$$

is valid. In the abbreviated form this implication can be written as

$$(a \leq b \vee a, b \in H) \wedge t(a) \sqsubset c \wedge a, t(b) \in X \longrightarrow c \in X. \quad (19)$$

Let $C_H(X)$ denotes the set of all $c \in G$ for which there exist $a, b \in G$ and $t \in T_n(G)$ such that the premise of (19) is satisfied. Further, let

$$C_H[X] = \bigcup_{m=0}^{\infty} \overset{m}{C}_H(X),$$

where $\overset{0}{C}_H(X) = X$, $\overset{m+1}{C}_H(X) = C_H(\overset{m}{C}_H(X))$ for every $m = 0, 1, 2, \dots$

By induction we can prove that $g \in \overset{m}{C}_H(X)$ if and only if the following system of conditions is fulfilled:

$$\left. \begin{aligned} & (a_1 = b_1[R_1 a \dots R_n a] \vee a_1, b_1 \in H) \wedge t_1(a_1)[R_1 g \dots R_n g] = t_1(a_1) \\ & \bigwedge_{i=1}^{2^{m-1}-1} \left(\begin{aligned} & a_{2i} = b_{2i}[R_1 a_{2i} \dots R_n a_{2i}] \vee a_{2i}, b_{2i} \in H, \\ & t_{2i}(a_{2i})[R_1 a_i \dots R_n a_i] = t_{2i}(a_{2i}), \\ & a_{2i+1} = b_{2i+1}[R_1 a_{2i+1} \dots R_n a_{2i+1}] \vee a_{2i+1}, b_{2i+1} \in H, \\ & t_{2i+1}(a_{2i+1})[R_1 t_i(b_i) \dots R_n t_i(b_i)] = t_{2i+1}(a_{2i+1}) \end{aligned} \right) \\ & \bigwedge_{i=2^{m-1}}^{2^m-1} (a_i \in X \wedge t_i(b_i) \in X) \end{aligned} \right\} \quad (20)$$

where $a_k, b_k \in G$, $t_k \in T_n(G)$.

In the sequel, the system of conditions (20) will be denoted by $\mathfrak{M}_H(X, m, g)$.

Theorem 2. *Let $\mathcal{G}$ be a functional Menger system of rank n . A nonempty subset H of G is a stabilizer of $\mathcal{G}$ if and only if it is a quasi-stable l -unitary normal v -complex such that $R_i H \subset H$ for every $i \in \overline{1, n}$ and*

$$x = y[R_1 x \dots R_n x] \in C_H[H] \wedge u[\bar{w} | i y] \in H \longrightarrow u[\bar{w} | i x] \in H \quad (21)$$

for all $x, y, u \in G$, $\bar{w} \in G^n$, $i \in \overline{1, n}$, where the symbol $u[\bar{w} | i]$ may be empty.

Proof. Necessity. Let H_Φ^a be the stabilizer of a point a in a functional Menger system $(\Phi, \mathcal{O}, \mathcal{R}_1, \dots, \mathcal{R}_n)$ of n -place functions. Obviously it is a quasi-stable l -unitary normal v -complex of $(\Phi, \mathcal{O}, \mathcal{R}_1, \dots, \mathcal{R}_n)$. If $f \in H_\Phi^a$, then $f(a, \dots, a) = a$, whence $\mathcal{R}_i f(a, \dots, a) = a$, i.e., $\mathcal{R}_i f \in H_\Phi^a$. So, $\mathcal{R}_i H_\Phi^a \subset H_\Phi^a$ for every $i \in \overline{1, n}$.

To prove (21), we shall consider $f = g[\mathcal{R}_1 f \dots \mathcal{R}_n f] \in C_{H_\Phi^a}[\mathcal{R}_1 f \dots \mathcal{R}_n f]$ and $\alpha[\bar{w} | i g] \in H_\Phi^a$ for some $f, g, \alpha \in \Phi$, $\bar{w} \in \Phi^n$, $i \in \overline{1, n}$, where $C_{H_\Phi^a}[\mathcal{R}_1 f \dots \mathcal{R}_n f] = \bigcup_{m=0}^{\infty} \overset{m}{C}_{H_\Phi^a}(\mathcal{R}_1 f \dots \mathcal{R}_n f)$. Then $f = g[\mathcal{R}_1 f \dots \mathcal{R}_n f] \in \overset{m}{C}_{H_\Phi^a}(\mathcal{R}_1 f \dots \mathcal{R}_n f)$ for some $m \in \mathbb{N}$. But,

as it is easily to see by induction, $\varphi \in \overset{m}{C}_{H_\Phi^a}(\mathcal{R}_1 f \dots \mathcal{R}_n f)$ implies $(a, \dots, a) \in \text{pr}_1 \varphi$. The above means that $(a, \dots, a) \in \text{pr}_1 f$ and $(a, \dots, a) \in \text{pr}_1 g[\mathcal{R}_1 f \dots \mathcal{R}_n f]$. Therefore

$$\begin{aligned} f(a, \dots, a) &= g[\mathcal{R}_1 f \dots \mathcal{R}_n f](a, \dots, a) \\ &= g(\mathcal{R}_1 f(a, \dots, a), \dots, \mathcal{R}_n f(a, \dots, a)) = g(a, \dots, a). \end{aligned}$$

Moreover, for $\alpha[\bar{w} | i g] \in H_\Phi^a$ we have

$$\begin{aligned} a &= \alpha[\bar{w} | i g](a, \dots, a) = \alpha(\bar{w}(a, \dots, a) | i g(a, \dots, a)) \\ &= \alpha(\bar{w}(a, \dots, a) | i f(a, \dots, a)) = \alpha[\bar{w} | i f](a, \dots, a), \end{aligned}$$

where $\bar{\omega}(a, \dots, a)$ denotes $\omega_1(a, \dots, a), \dots, \omega_n(a, \dots, a)$. So, $\alpha[\bar{\omega}|_i f] \in H_\Phi^a$, which completes the proof of (21).

Sufficiency. Let H satisfy all the conditions of the theorem. We prove that it satisfies also all the conditions of Theorem 1. Since by the assumption H is a quasi-stable l -unitary normal v -complex such that $R_i H \subset H$ for every $i \in \overline{1, n}$, we must prove that it satisfies the conditions (5), (6), (7) and $R_i U \subset H$, $R_i(G \setminus U) \subset G \setminus U$ for some $U \subset G$ containing H and all $i \in \overline{1, n}$.

First we prove that H satisfies the condition (8). Indeed, if $x \leq y$ and $x \in H$, then $y[R_1 x \dots R_n x] = x \in H$ and $R_i x \in H$, $i \in \overline{1, n}$. As H is an l -unitary normal v -complex, H is a v -unitary subset. Therefore $y \in H$, which completes the proof of (8).

Now let $U = C_H[H]$. Clearly $H \subset U$. Since $C_H[H] = \bigcup_{m=0}^{\infty} C_H^m(H)$, to prove that $R_i U \subset H$ for every $i \in \overline{1, n}$, it is sufficient to show that for every $m \in \mathbb{N}$ holds the inclusion

$$R_i(C_H^m(H)) \subset H. \quad (22)$$

For $m = 0$ it is obvious because $C_H^0(H) = H$ and $R_i H \subset H$ for all $i \in \overline{1, n}$. Suppose that (22) is valid for some $k \in \mathbb{N}$. We prove that it is valid for $k + 1$. Let $g \in C_H^{k+1}(H)$. Then

$$(a \leq b \vee a, b \in H) \wedge t(a) \sqsubset g \wedge a, t(b) \in C_H^k(H)$$

for some $a, b \in G$, $t \in T_n(G)$. From $a, t(b) \in C_H^k(H)$, according to our supposition, we get $R_i a, R_i t(b) \in H$. If $a \leq b$ and $a \in H$, then, by (8), we have $b \in H$. Thus $a, b, R_i t(b) \in H$. Whence $R_i t(a) \in H$ because H is a normal v -complex. The condition $t(a) \sqsubset g$ implies $R_i t(a) \leq R_i g$, which, by (8), gives $R_i g \in H$. In a similar way $a, b \in H$ and $t(a) \sqsubset g$ proves $R_i g \in H$. Thus we have shown that $R_i(C_H^{k+1}(H)) \subset H$. So, the inclusion (22) is valid for $m = k + 1$ and consequently for every $m \in \mathbb{N}$. Therefore $R_i U \subset H$ for every $i \in \overline{1, n}$.

To prove the inclusion $R_i(G \setminus U) \subset G \setminus U$ observe that it is equivalent to the condition

$$(\forall g \in G) (g \in G \setminus U \longrightarrow R_i g \in G \setminus U),$$

which can be written in the form

$$(\forall g \in G) (R_i g \in U \longrightarrow g \in U). \quad (23)$$

In the case $U = C_H[H]$ the last condition means that

$$(\forall g \in G)(\forall m \in \mathbb{N}) \left(R_i g \in C_H^m(H) \longrightarrow (\exists n \in \mathbb{N}) g \in C_H^n(H) \right). \quad (24)$$

Let $R_i g \in C_H^m(H)$ for some $g \in G$ and $m \in \mathbb{N}$. Considering $R_i g = R_i(R_i g)$ and (22), we conclude $R_i g \in H$. Thus $R_i g \leq R_i g$, $R_i g \sqsubset g$ and $R_i g \in H$.

Therefore $g \in C_H(H)$, i.e. $g \in \overset{n}{C}_H(H)$ for some $n \in \mathbb{N}$. This proves (24). So, $R_i(G \setminus U) \subset G \setminus U$ for $U = C_H[H]$ and $i \in \overline{1, n}$.

To prove (5) observe that for $x, y \in H$ and $t(x) \in U = C_H[H]$, just proved inclusion implies $R_i t(x) \in R_i U \subset H$. Thus $x, y, R_i t(x) \in H$. But H is a normal v -complex, hence $R_i t(y) \in H$. So, for $U = C[H]$, the condition (5) is satisfied. Also (6) is valid because for $U = C_H[H]$ it coincides with (21). Since the subset $C_H[H]$ is C_H -closed, the condition (7) is valid too. This means that H satisfies all the conditions of Theorem 1. Hence, H is a stabilizer of a functional Menger system $\mathcal{G}$. $\square$

As it is not difficult to see, the condition (21) is equivalent to the system of conditions $(A'_m)_{m \in \mathbb{N}}$, where

$$A'_m : x = y[R_1 x \dots R_n x] \wedge x \in \overset{m}{C}_H(H) \wedge u[\bar{w} | _i y] \in H \longrightarrow u[\bar{w} | _i x] \in H$$

for all $x, y, u \in G$, $\bar{w} \in G^m$, $i \in \overline{1, n}$. Since, by (20), the condition A'_m is equivalent to

$$A_m : x = y[R_1 x \dots R_n x] \wedge \mathfrak{M}_H(H, m, x) \wedge u[\bar{w} | _i y] \in H \longrightarrow u[\bar{w} | _i x] \in H,$$

the last theorem can be written in the form:

Theorem 3. *A nonempty subset H of G is a stabilizer of a functional Menger system $\mathcal{G}$ of rank n if and only if it is a quasi-stable l -unitary normal v -complex such that $R_i H \subset H$ for every $i \in \overline{1, n}$ and the system of conditions $(A_m)_{m \in \mathbb{N}}$ is satisfied.*

Now we shall characterize stabilizers of functional Menger λ -algebras.

Theorem 4. *A nonempty subset H of G is a stabilizer of a functional Menger λ -algebra $\mathcal{G}_\lambda = (G, o, \lambda, R_1, \dots, R_n)$ of rank n if and only if*

- 1) *it is a quasi-stable, λ -stable and v -unitary subset of G ,*
- 2) *there exists a subset U of G such that $H \subset U$, $R_i U \subset H$ and $R_i(G \setminus U) \subset G \setminus U$ for every $i \in \overline{1, n}$,*
- 3) *the following two implications*

$$x \in U \wedge y \in H \longrightarrow y[R_1 x \dots R_n x] \in H, \quad (25)$$

$$x \in U \wedge y \in U \longrightarrow y[R_1 x \dots R_n x] \in U \quad (26)$$

are valid for all $x, y \in G$.

Proof. The proof of the necessity of the conditions of the theorem is similar to the proof of the necessity of the conditions of Theorem 1. So, we prove only their sufficiency.

Let all these conditions be satisfied. First we shall show the implication

$$x \lambda y \in U \wedge u[\bar{w} | _i (y \lambda z)] \in U \longrightarrow u[\bar{w} | _i (x \lambda y \lambda z)] \in U \quad (27)$$

for $x, y, z \in G$ and $i \in \overline{1, n}$, where the symbol $u[\bar{w} |_i]$ may be empty. For this suppose that the premise of (27) is satisfied. Then, according to (26), we have

$$u[\bar{w} |_i(y \wedge z)][R_1(x \wedge y) \dots R_n(x \wedge y)] \in U.$$

Whence, by **A**₃, we obtain $u[\bar{w} |_i(y \wedge z)[R_1(x \wedge y) \dots R_n(x \wedge y)]] \in U$. From this, applying **A**₈, we conclude $u[\bar{w} |_i(z \wedge y[R_1(x \wedge y) \dots R_n(x \wedge y)])] \in U$, which, by **A**₉, implies $u[\bar{w} |_i(x \wedge y \wedge z)] \in U$. This completes the proof of (27).

Further, in the same way as in the proof of Theorem 1, we can show that the conditions (8), (9), (10) are satisfied and $G \setminus U$ is an l -ideal. Next, we consider the relation:

$$\mathcal{E}[U] = \{(g_1, g_2) \in G \times G \mid g_1 \wedge g_2 \in U \vee g_1, g_2 \in U'\}, \quad (28)$$

where $U' = G \setminus U$. The reflexivity and symmetry of this relation are obvious. Let $(g_1, g_2), (g_2, g_3) \in \mathcal{E}[U]$. If $g_1, g_2, g_3 \in U'$, then evidently $(g_1, g_3) \in \mathcal{E}[U]$. If $g_1 \wedge g_2, g_2 \wedge g_3 \in U$, then, according to (27), we have $g_1 \wedge g_2 \wedge g_3 \in U$. Whence, in view of (9) and the fact that $g_1 \wedge g_2 \wedge g_3 \leq g_1 \wedge g_3$, $\zeta \subset \chi$ and $g_1 \wedge g_2 \wedge g_3 \sqsubset g_1 \wedge g_3$, we conclude $g_1 \wedge g_3 \in U$, i.e., $(g_1, g_3) \in \mathcal{E}[U]$. So, $\mathcal{E}[U]$ is also transitive. Thus $\mathcal{E}[U]$ is an equivalence relation.

It is v -regular too. Indeed, if $(g_1, g_2) \in \mathcal{E}[U]$, then $g_1, g_2 \in U'$ or $g_1 \wedge g_2 \in U$. Since U' is an l -ideal, in the case $g_1, g_2 \in U'$ we have $u[\bar{w} |_i g_1], u[\bar{w} |_i g_2] \in U'$, i.e.,

$$u[\bar{w} |_i g_1] \equiv u[\bar{w} |_i g_2](\mathcal{E}[U]). \quad (29)$$

In the case when $g_1 \wedge g_2 \in U$ and $u[\bar{w} |_i g_1], u[\bar{w} |_i g_2] \in U'$, (29) is satisfied too. In the case $g_1 \wedge g_2 \in U$, $u[\bar{w} |_i g_1] \in U$, according to (26), we have

$$u[\bar{w} |_i g_1][R_1(g_1 \wedge g_2) \dots R_n(g_1 \wedge g_2)] \in U,$$

whence, applying **A**₃, we obtain

$$u[\bar{w} |_i g_1[R_1(g_1 \wedge g_2) \dots R_n(g_1 \wedge g_2)]] \in U,$$

which, by **A**₉, implies $u[\bar{w} |_i(g_1 \wedge g_2)] \in U$. This, in view of $u[\bar{w} |_i(g_1 \wedge g_2)] \sqsubset u[\bar{w} |_i g_2]$ and (9), gives $u[\bar{w} |_i g_2] \in U$.

Similarly we can prove that $g_1 \wedge g_2 \in U$ and $u[\bar{w} |_i g_2] \in U$ imply $u[\bar{w} |_i g_1] \in U$. Therefore $u[\bar{w} |_i g_1]$ and $u[\bar{w} |_i g_2]$ belong or do not belong to U simultaneously. Let $u[\bar{w} |_i g_1], u[\bar{w} |_i g_2] \in U$. Since

$$u[\bar{w} |_i(g_1 \wedge g_2)] \leq u[\bar{w} |_i g_j]$$

for $j = 1, 2$, from the above we obtain

$$u[\bar{w} |_i(g_1 \wedge g_2)] \sqsubset u[\bar{w} |_i g_1] \wedge u[\bar{w} |_i g_2],$$

which, after application of (9), gives $u[\bar{w} |_i(g_1 \wedge g_2)] \in U$. Thus (29) is satisfied in any case. So, the relation $\mathcal{E}[U]$ is i -regular for every $i \in \overline{1, n}$, i.e., it is v -regular.

H is an equivalence class of $\mathcal{E}[U]$. Indeed, if $g_1, g_2 \in H$, then $g_1 \wedge g_2 \in H$ by the $\wedge$ -stability of H . Consequently $g_1 \wedge g_2 \in U$, i.e., $g_1 \equiv g_2(\mathcal{E}[U])$. On the other hand, if $g_1 \equiv g_2(\mathcal{E}[U])$ and $g_1 \in H$, then $g_1 \wedge g_2 \in U$, whence, by (25), we have

$$g_1[R_1(g_1 \wedge g_2) \dots R_n(g_1 \wedge g_2)] \in H.$$

From this, applying **A**₉, we deduce $g_1 \wedge g_2 \in H$, which, in view of $g_1 \wedge g_2 \leq g_2$ and (8), implies $g_2 \in H$. So, H is an equivalence class of $\mathcal{E}[U]$.

Consider the simplest representation $P_{(\mathcal{E}[U], U')}$ of (G, o) induced by the pair $(\mathcal{E}[U], U')$ (see the section 2). We shall prove that this representation satisfies the following two identities:

$$P_{(\mathcal{E}[U], U')}(g_1 \wedge g_2) = P_{(\mathcal{E}[U], U')}(g_1) \cap P_{(\mathcal{E}[U], U')}(g_2), \quad (30)$$

$$P_{(\mathcal{E}[U], U')}(R_i g) = \mathcal{R}_i P_{(\mathcal{E}[U], U')}(g), \quad (31)$$

where $g, g_1, g_2 \in G$ and $i \in \overline{1, n}$.

Let $P_{(\mathcal{E}[U], U')}(g_1 \wedge g_2)(a_1, \dots, a_n) = b$, where H_{a_i}, H_b are equivalence classes of $\mathcal{E}[U]$ containing elements $x_i, y \in U$, respectively. Thus $(g_1 \wedge g_2)[x_1 \dots x_n] \equiv y(\mathcal{E}[U])$, whence

$$y \wedge g_1[x_1 \dots x_n] \wedge g_2[x_1 \dots x_n] \in U, \quad (32)$$

according to **A**₁₀. Consequently, $y \wedge g_1[x_1 \dots x_n] \in U$ and $y \wedge g_2[x_1 \dots x_n] \in U$, i.e., $g_1[x_1 \dots x_n] \equiv y(\mathcal{E}[U])$ and $g_2[x_1 \dots x_n] \equiv y(\mathcal{E}[U])$. Therefore

$$P_{(\mathcal{E}[U], U')}(g_1)(a_1, \dots, a_n) = b \quad \text{and} \quad P_{(\mathcal{E}[U], U')}(g_2)(a_1, \dots, a_n) = b. \quad (33)$$

This proves the inclusion $P_{(\mathcal{E}[U], U')}(g_1 \wedge g_2) \subset P_{(\mathcal{E}[U], U')}(g_1) \cap P_{(\mathcal{E}[U], U')}(g_2)$.

To prove the inverse inclusion assume (33). Then $g_1[x_1 \dots x_n] \equiv y(\mathcal{E}[U])$ and $g_2[x_1 \dots x_n] \equiv y(\mathcal{E}[U])$, i.e., $y \wedge g_1[x_1 \dots x_n] \in U$ and $y \wedge g_2[x_1 \dots x_n] \in U$. From this, according to (27), we obtain (32), whence, as it was shown above, we have $P_{(\mathcal{E}[U], U')}(g_1 \wedge g_2)(a_1, \dots, a_n) = b$. So, $P_{(\mathcal{E}[U], U')}(g_1) \cap P_{(\mathcal{E}[U], U')}(g_2) \subset P_{(\mathcal{E}[U], U')}(g_1 \wedge g_2)$. This completes the proof of (30).

Further, using the same method as in the proof of the condition (17), we can prove that

$$\text{pr}_1 P_{(\mathcal{E}[U], U')}(R_i g) = \text{pr}_1 P_{(\mathcal{E}[U], U')}(g) = \text{pr}_1 \mathcal{R}_i P_{(\mathcal{E}[U], U')}(g)$$

for every $g \in G$. Now if $(a_1, \dots, a_n) \in \text{pr}_1 P_{(\mathcal{E}[U], U')}(R_i g)$, then $(R_i g)[x_1 \dots x_n] \in U$ for all x_i from the class H_{a_i} , $i \in \overline{1, n}$. Since $(R_i g)[x_1 \dots x_n] \leq x_i$ implies $(R_i g)[x_1 \dots x_n] \wedge x_i = (R_i g)[x_1 \dots x_n]$, the above gives $(R_i g)[x_1 \dots x_n] \wedge x_i \in U$. Hence $(R_i g)[x_1 \dots x_n] \equiv x_i(\mathcal{E}[U])$. Therefore,

$$P_{(\mathcal{E}[U], U')}(R_i g)(a_1, \dots, a_n) = a_i$$

for all $i \in \overline{1, n}$. This proves (31). So, the simplest representation $P_{(\mathcal{E}[U], U')}$ of (G, o) is also the simplest representation of a functional Menger $\wedge$ -algebra $(G, o, \wedge, R_1, \dots, R_n)$. Moreover, analogously as (18), we can prove that

$$g \in H \longleftrightarrow P_{(\mathcal{E}[U], U')}(g)(a, \dots, a) = a,$$

where a is an elements used as index of the $\mathcal{E}[U]$ -class H .

Since the algebra $\mathcal{G}_\lambda$ satisfies the axioms $\mathbf{A}_1 - \mathbf{A}_{10}$, from [1] it follows that there exists an isomorphism of $\mathcal{G}$ onto some functional Menger $\cap$ -algebra of n -place functions. Denote this isomorphism by P_1 and consider the representation $P = P_1 + P_{(\mathcal{E}[U], U')}$. It is clear that P is a faithful representation of $\mathcal{G}$ by n -place functions and $H = H_P^a$. $\square$

Theorem 5. *A nonempty subset H of a functional Menger λ -algebra $\mathcal{G}_\lambda$ of rank n is its stabilizer if and only if it is stable, λ -stable and v -unitary subset of G such that $R_i H \subset H$ for every $i \in \overline{1, n}$.*

Proof. The necessity of these conditions is obvious, therefore we shall prove only their sufficiency.

Let H satisfies all these conditions. Then $\zeta(H) \subset H$ which is equivalent to (8).² Indeed, for $x \in H$ and $x \leq y$ we have $y[R_1 x \dots R_n x] = x \in H$ and $R_i x \in H$ for ever $i \in \overline{1, n}$, whence, according to the v -unitarity of H , we obtain $y \in H$. This proves (8).

Using this condition we shall prove that $U_0 = \chi(H)$ and H satisfy all conditions of Theorem 4. The stability of H implies its quasi-stability. Because χ is a quasi-order we have also $H \subset \chi(H) = U_0$. Moreover, for every $x \in U_0$ there exists $h \in H$ such that $(h, x) \in \chi$, i.e., $R_i h \leq R_i x$ for every $i \in \overline{1, n}$. Since $R_i h \in H$ for every $h \in H$ and $i \in \overline{1, n}$, the above, according to (8), implies $R_i x \in H$ for $i \in \overline{1, n}$. Therefore $R_i U_0 \subset H$ for every $i \in \overline{1, n}$.

The inclusion $R_i(G \setminus U_0) \subset G \setminus U_0$, where $i \in \overline{1, n}$, is equivalent to the implication $(\forall x \in G)(x \in G \setminus U_0 \longrightarrow R_i x \in G \setminus U_0)$, which, by contraposition, means that

$$R_i x \in U_0 \longrightarrow x \in U_0$$

for every $x \in G$. But $U_0 = \chi(H)$, so, for every $x \in G$ and every $h \in H$

$$(h, R_i x) \in \chi \longrightarrow x \in \chi(H). \quad (34)$$

Let $(h, R_i x) \in \chi$ for some $x \in G$ and $h \in H$. Then

$$H \ni h = h[R_1 R_i x \dots R_n R_i x] = h[R_1 x \dots R_n x],$$

whence $(h, x) \in \chi$. Therefore $x \in \chi(H)$. This means that the implication (34) is valid. So, $R_i(G \setminus U_0) \subset G \setminus U_0$ for every $i \in \overline{1, n}$. In this way we have proved that U_0 and H satisfy the first two conditions of Theorem 4.

To prove that U_0 and H satisfy the third condition of this theorem, we must show the following two implications:

$$x \in U_0 \wedge y \in H \longrightarrow y[R_1 x \dots R_n x] \in H, \quad (35)$$

$$x \in U_0 \wedge y \in U_0 \longrightarrow y[R_1 x \dots R_n x] \in U_0. \quad (36)$$

² Remind that for any relation $\rho \subset X \times Y$ and any subset A of X by $\rho(A)$ is denoted the set $\{y \in Y \mid (\exists x \in A)(x, y) \in \rho\}$.

Let $x \in U_0 = \chi(H)$ and $y \in H$. Then, $(h, x) \in \chi$, i.e., $h = h[R_1x \dots R_nx]$ for some $h \in H$. Since from $h \in H$ it follows $R_ih \in H$ for all $i \in \overline{1, n}$, we have $y, R_ih \in H$, $i \in \overline{1, n}$. This, by the stability of H , implies $y[R_1h \dots R_nh] \in H$. Whence, in view of

$$\begin{aligned} y[R_1h \dots R_nh] &= y[R_1h[R_1x \dots R_nx] \dots R_nh[R_1x \dots R_nx]] \\ &\stackrel{\mathbf{A}_4}{=} y[(R_1h)[R_1x \dots R_nx] \dots (R_nh)[R_1x \dots R_nx]] \\ &\stackrel{\mathbf{A}_1}{=} y[R_1h \dots R_nh][R_1x \dots R_nx] \\ &\stackrel{\mathbf{A}_5}{=} y[R_1x \dots R_nx][R_1h \dots R_nh], \end{aligned}$$

we conclude $y[R_1x \dots R_nx][R_1h \dots R_nh] \in H$. But H is v -unitary and $R_ih \in H$ for every $i \in \overline{1, n}$, so, $y[R_1x \dots R_nx] \in H$. This proves (35).

Now let $x, y \in U_0 = \chi(H)$. Then there exist $a, b \in H$ such that $(a, x) \in \chi$ and $(b, y) \in \chi$, i.e., $a = a[R_1x \dots R_nx]$ and $b = b[R_1y \dots R_ny]$. Because $b \in H$ implies $R_ib \in H$ for every $i \in \overline{1, n}$, we have $a, R_ib \in H$ for all $i \in \overline{1, n}$. Whence, according to the stability of H , we get $a[R_1b \dots R_nb] \in H$. Moreover,

$$\begin{aligned} a[R_1b \dots R_nb] &= a[R_1x \dots R_nx][R_1b \dots R_nb] \\ &\stackrel{\mathbf{A}_5}{=} a[R_1b \dots R_nb][R_1x \dots R_nx] \\ &= a[R_1b[R_1y \dots R_ny] \dots R_nb[R_1y \dots R_ny]] \\ &\stackrel{\mathbf{A}_4}{=} a[(R_1b)[R_1y \dots R_ny] \dots (R_nb)[R_1y \dots R_ny]][R_1x \dots R_nx] \\ &\stackrel{\mathbf{A}_1}{=} a[R_1b \dots R_nb][R_1y \dots R_ny][R_1x \dots R_nx] \\ &\stackrel{\mathbf{A}_1}{=} a[R_1b \dots R_nb][(R_1y)[R_1x \dots R_nx] \dots (R_ny)[R_1x \dots R_nx]] \\ &\stackrel{\mathbf{A}_4}{=} a[R_1b \dots R_nb][R_1(y[R_1x \dots R_nx]) \dots R_n(y[R_1x \dots R_nx])]. \end{aligned}$$

Thus, $(a[R_1b \dots R_nb], y[R_1x \dots R_nx]) \in \chi$, i.e., $y[R_1x \dots R_nx] \in U_0 = \chi(H)$. This completes the proof of (36).

In this way we have shown that U_0 and H satisfy all the conditions of Theorem 4. Therefore, H is the stabilizer of a functional Menger $\wedge$ -algebra $\mathcal{G}_\wedge$. $\square$

References

- [1] Dudek W.A., Trokhimenko V.S.: *Functional Menger $\mathcal{P}$ -algebras*, Commun. Algebra, 30 (2002), 5921 – 5931.
- [2] Dudek W.A., Trokhimenko V.S.: *Representations of Menger $(2, n)$ -semi-groups by multiplace functions*, Commun. Algebra, 34 (2006), 259 – 274.
- [3] Dudek W.A., Trokhimenko V.S.: *Menger algebras of multiplace functions*, (Russian), Centrul Ed. USM, Chişinău 2006.

- [4] Schein B. M.: *Stationary subsets, stabilizers and transitive representations of semigroups*, Dissertationes Math., 77 (1970), 3 – 36.
- [5] Schein B. M.: *Lectures on semigroups of transformations*, Amer. Math. Soc. Translat. (2), 113 (1979), 123 – 181.
- [6] Schein B. M., Trokhimenko V. S.: *Algebras of multiplace functions*, Semigroup Forum, 17 (1979), 1 – 64.
- [7] Trokhimenko V. S.: *Menger's function systems*, (Russian), Izv. Vyssh. Uchebn. Zaved. Matematika, 11 (1973), 71 – 78.
- [8] Trokhimenko V. S.: *Stabilizers of ordered Menger algebras*, (Russian), Izv. Vyssh. Uchebn. Zaved. Matematika, 6 (1986), 74 – 76.
- [9] Trokhimenko V. S.: *Stationary subsets and stabilizers of restrictive Menger $\mathcal{P}$ -algebras of multiplace functions*, Algebra Universalis, 44 (2000), 129 – 142.

Dudek W. A.
 Institute of Mathematics and
 Computer Science
 Wroclaw University of Technology
 50-370 Wroclaw
 Poland
 E-mail: dudek@im.pwr.wroc.pl

Trokhimenko V. S.
 Department of Mathematics
 Pedagogical University
 21100 Vinnitsa
 Ukraine
 E-mail: vtrokhim@sovamua.com